

Acetaminophen (Tylenol) Dosing Information

Give every 4-6 hours, as needed, and not more than five times in 24 hours unless directed by a health care professional.

| Weight | Age | Infant Oral Suspension: Concentration 5 mL = 160mg | Children's Suspension 1 tsp (5 mL) = 160 mg | Children's Tablets 1 tablet = 80mg | Junior Strength 1 tablet = 160 mg |
|-----------------|---|---|---|---|--|
| 6-11 pounds | 0-3 months only to be given if directed by a health care professional (see above) | | | | |
| 12-17 pounds | 4-11 months | 2.5mL | 1/2 teaspoon (80 mg) | | |
| 18-23 pounds | 12-23 months | 3.75mL | 3/4 teaspoon (120 mg) | | |
| 24-35 pounds | 2-3 years | 5mL | 1 teaspoon (160 mg) | 2 tablets | |
| 36-47 pounds | 4-5 years | | 1 1/2 teaspoons (240 mg) | 3 tablets | |
| 48-59 pounds | 6-8 years | | 2 teaspoons (320 mg) | 4 tablets | 2 tablets |
| 60-71 pounds | 9-10 years | | 2 1/2 teaspoons (400 mg) | 5 tablets | 2.5 tablets |
| 72-95 pounds | 11 years | | 3 teaspoons (480 mg) | 6 tablets | 3 tablets |

